

Cardiac failure congestive	1	0
Cardiorespiratory arrest	1	1
Chronic obstructive pulmonary disease	1	
Death	0	1
Dementia		
Emphysematous cholecystitis	1	0
Hemorrhagic stroke		1
Hypertensive heart disease	1	0
Lung cancer metastatic	1	0
Metastases to liver	0	
Missing	0	1
Multiple organ dysfunction syndrome	0	
Myocardial infarction	0	2
Overdose	0	1
Pneumonia	0	2
Sepsis	1	0
Septic shock	1	0
Shigella sepsis		0
Unevaluable event	1	0

Table S4 | Causes of Death from Dose 1 to Unblinding (Safety Population, ≥ 16 Years Old). a.
Multiple causes of death could be reported for each participant. There were no deaths among 12–15-year-old participants.

EuropeReloaded.com

It is linked to below, but it's worth following Dr. David White as he walks us through some data on the original Pfizer trials. What was interesting to us is how they mask

Here's Russell's own transcribed version of the tweet by Dr. White

7:06 AM · Jun 29, 2024



757



Reply



Share this post

[Read 17 replies](#)

AUSSIE17

It's important for people to understand how Pfizer manipulated the categorization of deaths in their original trial, which led to politicians using the "safe and effective" narrative. Everyone should carefully watch this, as it not only raises concerns about

months, as ~~published~~ in the New England Journal of Medicine. The original trial showed that **deaths were about 7% higher in the vaccine group**, with 15 deaths, compared to 14 deaths in the placebo group. However, many “fact-checkers” and pharma shills “scientists” claim that this difference is ***not statistically significant***.

This is intriguing because, according to a Lancet paper, there were 44,000 more deaths in England in 2022, driven by a **sharp increase in cardiovascular deaths** among the middle-aged, which rose by 33%. **He wonders if cardiovascular deaths in the trial's vaccine group were also 33% higher.** This question is important because it seeks to determine if the higher death rate in the vaccinated group is related to the rise in cardiovascular deaths seen in the general population.

He walks us through four mysterious “causes of deaths” categorized in the paper:

Pneumonia	0	2
Sepsis	1	0
Septic shock	1	0
Shigella sepsis	0	0
Unevaluable event	1	0

Table S4 | Causes of Death from Dose 1 to Unblinding (Safety Population, ≥16 Years Old). a.
Multiple causes of death could be reported for each participant. There were no deaths among 12–15-year-old participants.

- An “Unevaluable Event” on a vaccinated death, which was found to be a **sudden cardiac death confirmed via autopsy**.
 - A “Missing” entry categorized as a “placebo” death, which was **later found to be a “vaccinated” death**.
 - “Emphysematous Cholecystitis,” which is a cardiovascular death but Pfizer said let’s put something that sounds really complicated and hope nobody knows **it’s a cardiovascular death**.
 - “Death” for participant #10841470. *First of all categorizing a “cause of death” as “Death”, are you kidding?* Anyway this participant **died after taking a Moderna mRNA vaccine, but was categorized as a placebo death**. I wrote about this issue [here](#):

Dr. White highlights that these figures, particularly the increased all-cause mortality in the vaccine group, **clearly point to safety signals**. He stresses that informed consent is a fundamental ethical principle in medicine, necessitating the open discussion of such safety concerns. It is crucial to ensure that participants are fully informed about the potential risks associated with the vaccine to maintain trust and transparency within the medical community.

Subscribe to Truth11.com

Receive Articles By Email • Support Independent Media

 **Subscribe now**

Support Truth11.com • Make A Donation

• Or Become A Monthly Subscriber

